

# The CANADIAN FIELD-NATURALIST

A JOURNAL OF FIELD BIOLOGY AND ECOLOGY

Promoting the study and conservation of northern biodiversity since 1880



Volume 133, Number 1 • January–March 2019



Ottawa Field-Naturalists' Club  
Club des naturalistes d'Ottawa

# The Ottawa Field-Naturalists' Club

FOUNDED 1863 (CURRENT INCORPORATION 1879)

## Patron

Her Excellency the Right Honourable Julie Payette, C.C., C.M.M., C.O.M., C.Q., C.D.  
Governor General of Canada

The objectives of this Club shall be to promote the appreciation, preservation, and conservation of Canada's natural heritage; to encourage investigation and publish the results of research in all fields of natural history and to diffuse information on these fields as widely as possible; to support and cooperate with organizations engaged in preserving, maintaining, or restoring environments of high quality for living things.

## Honorary Members

Ronald E. Bedford	Michael D. Cadman	J. Bruce Falls	Robert E. Lee	Allan H. Reddoch
Charles D. Bird	Paul M. Catling	Peter W. Hall	John McNeill	Joyce M. Reddoch
Fenja Brodo	Francis R. Cook	Christine Hanrahan	Theodore Mosquin	Dan Strickland
Irwin M. Brodo	Bruce Di Labio	C. Stuart Houston	Robert W. Nero	John B. Theberge
Daniel F. Brunton	Anthony J. Erskine	Ross A. Layberry	E. Franklin Pope	Sheila Thomson

## 2019 Board of Directors

<b>President:</b> Diane Lepage	Annie Bélair	Edward Farnworth	Dwayne Lepitzki	Henry Steger
<b>1st Vice-President:</b> Jakob Mueller	Fenja Brodo	Catherine Hessian	Bev McBride	Ken Young
<b>Recording Secretary:</b> Elizabeth Moore	Robert Cermak	Anouk Hoedeman	Gordon Robertson	Eleanor Zurbrigg
<b>Treasurer:</b> Ann Mackenzie	Owen Clarkin	Diane Kitching	Jeff Saarela	

To communicate with the Club, address postal correspondence to: The Ottawa Field-Naturalists' Club, P.O. Box 35069, Westgate P.O., Ottawa, ON, K1Z 1A2, or e-mail: [ofnc@ofnc.ca](mailto:ofnc@ofnc.ca). For information on club activities, go to [www.ofnc.ca](http://www.ofnc.ca).

## The Canadian Field-Naturalist

The Canadian Field-Naturalist is published quarterly by The Ottawa Field-Naturalists' Club. Opinions and ideas expressed in this journal do not necessarily reflect those of The Ottawa Field-Naturalists' Club or any other agency.

**Website:** [www.canadianfieldnaturalist.ca/index.php/cfn](http://www.canadianfieldnaturalist.ca/index.php/cfn)

**Editor-in-Chief:** Dr. Dwayne Lepitzki

**Copy Editors:** Sandra Garland and Dr. John Wilmshurst

**Book Review Editor:** Dr. Barry Cottam

**Subscription Manager:** Eleanor Zurbrigg

**Assistant Editor:** Dr. Amanda Martin

**Layout:** Robert Forsyth

**Online Journal Manager:** Dr. Bill Halliday

**Author Charges Manager:** Ken Young

<b>Associate Editors:</b>	Dr. Ron Brooks	Dr. Jennifer R. Foote	Dr. Donald F. McAlpine	Dr. Jeffery M. Saarela
	Dr. Carolyn Callaghan	Dr. Graham Forbes	Dr. Garth Mowat	David C. Seburn
	Dr. Paul M. Catling	Thomas S. Jung	Dr. Marty Obbard	Dr. Jeffrey H. Skevington
	Dr. François Chapleau			

**Chair, Publications Committee:** Dr. Jeffery M. Saarela

**All manuscripts intended for publication**—except Book Reviews—should be submitted through the online submission system at the CFN website: <http://www.canadianfieldnaturalist.ca/index.php/cfn/user>. Click the “New Submission” link on the right side of the webpage and follow the prompts. Authors must register for a cfn account at <http://www.canadianfieldnaturalist.ca/index.php/cfn/user/register> in order to submit a manuscript. Please contact the Online Journal Manager ([info@canadianfieldnaturalist.ca](mailto:info@canadianfieldnaturalist.ca)) if you have any questions or issues with the online submission process. In only rare, exceptional circumstances will submissions other than online be considered and, in these cases, authors must contact the Editor-in-Chief ([editor@canadianfieldnaturalist.ca](mailto:editor@canadianfieldnaturalist.ca)) prior to submission. Instructions for Authors are found at <http://www.canadianfieldnaturalist.ca/public/journals/1/CFNAuthorinstructions.pdf>.

**Book-review correspondence**, including arranging for delivery of review copies of books, should be sent to the Book Review Editor by e-mail: [b.cottam@rogers.com](mailto:b.cottam@rogers.com).

**Subscriptions and Membership:** Subscription rates for individuals are \$40 (online only), \$50 (print only), or \$60 (print + online). Libraries and other institutions may subscribe for \$120 (online only or print only) or \$180 (print + online). All foreign print subscribers and members (including USA) must add \$10 to cover postage. The Ottawa Field-Naturalists' Club annual membership fee of \$40 (individual), \$45 (family), or \$20 (student) includes an online subscription to *The Canadian Field-Naturalist*. Members can receive printed issues of CFN for an additional \$30 per volume (four issues). For further details, see <http://ofnc.ca/membership-and-donations>. The club's regional journal, *Trail & Landscape*, covers the Ottawa District and regional Club events and field trips. It is mailed to all club members. It is available to libraries at \$40 per year. Subscriptions, applications for membership, notices of changes of address, and undeliverable copies should be sent to [subscriptions@canadianfieldnaturalist.ca](mailto:subscriptions@canadianfieldnaturalist.ca) or mailed to: The Ottawa Field-Naturalists' Club, P.O. Box 35069, Westgate P.O., Ottawa, ON, K1Z 1A2 Canada. Canada Post Publications Mail Agreement number 40012317. Return postage guaranteed.

The Thomas H. Manning fund, a special fund of the OFNC, established in 2000 from the bequest of northern biologist Thomas H. Manning (1911–1998), provides financial assistance for the publication of papers in the CFN by independent (non-institutional) authors, with particular priority given to those addressing Arctic and boreal issues. Qualifying authors should make their application for assistance from the fund at the time of their initial submission.

**COVER:** Rock Ptarmigan (*Lagopus muta townsendi*) on Amchitka Island, Alaska, in late May to very early June. Male Rock Ptarmigan on mainland Alaska and northern Canada during this same period are mostly white, a difference that has been used to describe subspecies of Rock Ptarmigan. See the article in this issue by Clait Braun *et al.*, pages 49–55. Photo: Steve Ebbert, 10 June 2015.

## Monitoring Rock Ptarmigan (*Lagopus muta*) populations in the Western Aleutian Islands, Alaska

CLAIT E. BRAUN<sup>1,\*</sup>, WILLIAM P. TAYLOR<sup>2</sup>, STEVEN M. EBBERT<sup>3,4</sup>, and LISA M. SPITLER<sup>5</sup>

<sup>1</sup>Grouse Inc., 5572 North Ventana Vista Road, Tucson, Arizona 85750 USA

<sup>2</sup>12841 Nora Drive, Anchorage, Alaska 99515 USA

<sup>3</sup>Alaska Maritime National Wildlife Refuge, 95 Sterling Highway, Suite 101, Homer, Alaska 99603 USA

<sup>4</sup>Current Address: P.O. Box 457, Anchor Point, Alaska 99556 USA

<sup>5</sup>Alaska Maritime National Wildlife Refuge, Aleutian Islands Unit, P.O. Box 5251, Adak, Alaska 99546 USA

\*Corresponding author: sgwtp66@gmail.com

Braun, C.E., W.P. Taylor, S.M. Ebbert, and L.M. Spitler. 2019. Monitoring Rock Ptarmigan populations in the Western Aleutian Islands, Alaska. *Canadian Field-Naturalist* 133(1): 49–55. <https://doi.org/10.22621/cfn.v133i1.1948>

### Abstract

Knowledge of population fluctuations of Aleutian Islands Rock Ptarmigan (*Lagopus muta*) is limited because of isolation and access. We reviewed the available but limited data on ptarmigan counts on islands in North America and evaluated the use of point counts to estimate changes in apparent numbers of Rock Ptarmigan on three islands (Adak, Amchitka, and Attu) in the Western Aleutian Islands in Alaska. We developed a standardized protocol to count numbers of Rock Ptarmigan (males and females) seen and/or heard on 5-minute point counts at 0.8 km intervals along marked global positioning system routes on Adak (2015–2017), Amchitka (2015), and Attu (2015) islands. Apparent densities based on Rock Ptarmigan seen and/or heard at 98 stops on 10 routes varied and were highest (1.9 birds per stop in 2015, 1.4 in 2016, and 1.0 in 2017) on Adak, lower (0.4 birds per stop) on Amchitka, and lowest (0.0 birds per stop) on Attu in late May–early June 2015. These island populations represent three subspecies and unique conservation units. Continuation of point-count surveys of these three subspecies in future years will provide baseline data over time and lead to a better understanding of any fluctuations in and synchrony among Rock Ptarmigan populations on these islands. This information is necessary for both theoretical (how are ptarmigan breeding populations regulated on islands) and practical reasons (identifying the optimal period for possible translocation to islands where ptarmigan were extirpated by introduced Arctic Fox [*Vulpes lagopus*]).

Key words: Rock Ptarmigan; *Lagopus muta*; Adak; Amchitka; Attu; Aleutian Islands; point counts; Alaska; USA

### Introduction

Animal population fluctuations have long been of interest (Elton 1924), especially in insular areas that have no obvious corridors or where populations are otherwise isolated. Rock Ptarmigan (*Lagopus muta*) has a circumpolar distribution in northern latitudes with multiple subspecies: up to 14 in North America alone (AOU 1957; the last time subspecies were listed by the American Ornithologists' Union). Populations of Rock Ptarmigan occupy remote areas and their distribution can be highly fragmented including on islands. Thus, documentation of population fluctuations over time can be difficult. It is important to monitor the status and population changes of species, such as Rock Ptarmigan, and to investigate any underlying factors affecting long-term changes (Pedersen *et al.* 2005; Tesar *et al.* 2016). Measuring changes over time can be problematic in isolated areas such as in the Arctic and substantial efforts to learn how to effectively monitor population status of ptarmigan have

been made (Pelletier and Krebs 1997; Bart *et al.* 2011).

There is some evidence that Rock Ptarmigan are cyclic on islands (Iceland; Magnússon *et al.* 2004), but population trends are poorly documented in North America (Weeden 1965; Cotter 1999; Taylor 2013). Peaks in Rock Ptarmigan cycles may represent a 10-fold increase from lows as discussed by Holder and Montgomerie (1993: 15), who cited studies in Scotland (Watson 1965) and Canada (Cotter 1991). Grouse cycles may be correlated with changes in their predator numbers or parasites and not only immigration or emigration from the local population (Dobson and Hudson 1992; Hudson *et al.* 1992; Cattadori *et al.* 2005).

As many as seven to eight subspecies of Rock Ptarmigan have been described from the Aleutian Archipelago, Alaska (AOU 1957). This number has been condensed into four groups (*L. m. evermanni*, *L. m. townsendi*, *L. m. atkensis*, and *L. m. nelsoni*) of which *nelsoni* also occur on mainland Alaska to

A contribution towards the cost of this publication has been provided by the Thomas Manning Memorial Fund of the Ottawa Field-Naturalists' Club.

the east (Gibson and Kessel 1997; Montgomerie and Holder 2008). The *evermanni* subspecies occurs in the Near Islands (Attu and Agattu); *townsendi* occurs in the Rat Islands, including Amchitka and Kiska islands, while *atkensis* occurs in the Andreanof Islands group including Adak, Tanaga to Atka, and possibly other islands. We studied three subspecies (*atkensis*, *evermanni*, and *townsendi*), all of which are considered endemic groups or unique conservation units (Pruett *et al.* 2010).

Pelletier and Krebs (1997) tested line transect methods to estimate densities of breeding male ptarmigan and concluded they cannot be censused in small areas alone (size of their multiple study sites ranged from 3.0 to 13.5 km<sup>2</sup>) as the results were too variable. Others (Brodsky and Montgomerie 1987; Cotter 1999; Watson *et al.* 2000; Favaron *et al.* 2006; Pedersen *et al.* 2012) used methods such as marking and reobservation, point transects, and distance sampling to estimate changes in population size. Bart *et al.* (2011) experimented with use of helicopter and fixed-wing aircraft to survey ptarmigan (and other species) over large areas in northern Canada and Alaska. All of these methods are either labour intensive and/or costly.

The Breeding Bird Survey protocol was developed in the early 1960s to estimate population trends in bird populations over time across large areas and variable habitats in North America (Bystrak 1981; Robbins *et al.* 1986; Droege 1990). It initially used 3-minute count intervals along routes with 50 stops at 0.8 km

intervals with routes being surveyed once each year, during the breeding period. More recently 5-minute point counts have been used to better represent population indices of selected species (Ralph *et al.* 1995). All birds (males and females) both seen and heard are recorded at point-count stops.

Male Rock Ptarmigan during the breeding period (late April to early June) are conspicuous (they can range in colour from white to mottled shades of light brown to almost black [Attu] with white bellies and wings), perch up, and make flights from conspicuous sites while calling as they advertise and defend territories (Holder and Montgomerie 1993; Pelletier and Krebs 1997). Males in late May and early June can be solitary or paired with females, which are drab brown (in flight they have white wings) or mottled and inconspicuous in many, if not most, situations (some may be on nests).

Our objectives were to review the available literature on surveys of Rock Ptarmigan on three islands in the Western Aleutian Islands and develop and implement a 5-minute point count protocol to estimate trends in breeding populations of Rock Ptarmigan on Adak, Amchitka, and Attu islands in the Western Aleutian Islands (Figure 1).

### Study Area

We reviewed the published and other available literature on Rock Ptarmigan (no other species of ptarmigan occur on these islands) on Adak (51.883°N, 176.65°W), Amchitka (51.35°–51.65°N, 178.617°–




FIGURE 1. Aleutian Archipelago, Alaska showing Adak, Amchitka, and Attu islands.

179.483°E), and Attu (52.85°N, 173.183°E) in the Western Aleutian Islands, Alaska (Appendix S1). Areas surveyed on the three islands by us and others were similar low elevation sites (i.e., marine and stream terraces) adjacent to rarely-used trails (Amchitka and Attu) and occasionally used roads (Adak) that tended to follow coastal areas. The islands vary in size from ~300 km<sup>2</sup> (Amchitka) to 711 km<sup>2</sup> (Adak) and 894 km<sup>2</sup> (Attu). Adak is in the Andréanof group while Amchitka is in the Rat Island group and Attu is in the Near Islands. All are bounded by the North Pacific Ocean to the south and west and the Bering Sea to the north and east. The three islands are distant from each other with Amchitka being 301 km southwest of Adak that is 720 km east of Attu. There are no human residents on Amchitka and Attu and the resident population on Adak is variable and <100 people.

The geology of the three islands is complex with multiple inactive volcanos and volcanic flows as well as past glacial and marine erosion (Coats 1956; Fraser and Snyder 1959; Powers *et al.* 1960). Topography varies from gently sloping marine terraces to undulating tundra ranging to rugged mountains. We surveyed ptarmigan at an elevation of 10 to 300 m on all three islands. Lower and well-drained sites are occupied by grasses and sedges (*Calamagrostis* spp., *Leymus* spp., *Carex* spp.), and low-growing forbs including *Caltha* spp., *Ranunculus* spp., and *Lupinus* spp. with higher slopes dominated by crowberry (*Empetrum* spp.), *Empetrum-Cladonia* tundra, *Cladina* spp. lichens, and other mosses with some low-growing heather

(*Cassiope* spp., *Phyllodoce* spp.), willow (*Salix* spp.), and Kamchatka Rhododendron (*Rhododendron camtschaticum* Pallas) shrubs (Amundsen and Clebsch 1971; Everett 1971; White *et al.* 1977).

The climate on all three islands is moist marine with frequent high velocity winds, rain, and fog (Gates *et al.* 1971). Mean daily average temperatures vary seasonally ranging from 0.4°C in January to 11°C in August. Daily (3.9°C in all seasons) and seasonal (9.4°C) ranges in temperature are limited (Armstrong 1971, 1977). Wind speeds are highly variable, and the mean annual precipitation ranges from 83 to 139 cm, depending on the island, with June and July being the months with lowest precipitation (Weatherbase 2015).

## Methods

We established and conducted 5-minute point stations (protocol in Table 1) in 2015 following Ralph *et al.* (1995) at 0.8 km intervals along trails and roads on all three islands (dates in Table 2). All routes were conducted using an all-terrain vehicle. Starting points were at trail junctions or easily recognized local features and were recorded as global positioning system coordinates (on file with Alaska Maritime National Wildlife Refuge, Homer, Alaska, USA). Point-count routes were in areas where at least four stops at 0.8 km intervals could be established. There were four routes on Adak with from six to 17 stops, three routes on Amchitka with from four to 21 stops, and three routes with from four to seven stops on Attu (Table 2).

**TABLE 1.** Protocol for Rock Ptarmigan (*Lagopus muta*) surveys on Adak, Amchitka, and Attu islands, Western Aleutians, Alaska.

---



---

### 5-minute point counts

Count and record all Rock Ptarmigan seen (as male or female) or heard at each point stop. Rarely, the vehicle stopped near a ptarmigan which did not call or flush during the 5-minute count but flushed when the vehicle departed. These birds were included in the count.

Use of two counters is best (one front and one back). A central spot should be chosen.

Consolidate and record totals at end of each point count before returning to the vehicle.

Conduct point-count route prior to 10:00 AM.

Try to avoid high winds (>30 km per hr) and heavy rain.

### Adak (use GPS locations at >0.8 km points)

Old Loran Road (10 stops).

Finger Cove (six stops).

Navfac Creek to past Clam Cove (17 stops).

Lake Andrew (six stops).

### Amchitka (use GPS locations at >0.8 km points)

Jones Lake/Engineer Road (17 stops).

Charlie-Baker Taxiway south (four stops).

Infantry Road (21 stops).

Attu (use GPS locations at >0.8 km points)

Casco Cove to old airstrips (four stops).

Engineer Hill (top towards Peace monument) from Massacre Creek Beach Trail (seven stops).

Navytown (two stops) to Quonset Valley (four stops).

---



---

**TABLE 2.** Point-count (5-minute) surveys of Rock Ptarmigan (*Lagopus muta*; RP) on Adak, Amchitka, and Attu islands, Alaska, 2015–2017.

Island	Routes	Date	<i>n</i> points	RP seen/heard	Birds/Stop
Adak 2015	Finger Bay	29 May 2015	6	6	1.0
	Navfac Creek-Clam Lagoon	29 May 2015	17	12	0.7
	Old Loran Station Road	30 May 2015	10	49	4.9
	Andrew Lake	30 May 2015	6	6	1.0
Mean					1.9
Adak 2016	Finger Bay	18 May 2016	6	3	0.5
	Navfac Creek-Clam Lagoon	3 June 2016	17	16	1.4
	Old Loran Station Road	27 May 2016	10	10	2.1
	Andrew Lake	20 May 2016	6	10	1.7
Mean					1.4
Adak 2017	Finger Bay	1 June 2017	6	0	0.0
	Navfac Creek-Clam Lagoon	29 May 2017	17	16	0.9
	Old Loran Station Road	30 May 2017	10	16	1.6
	Andrew Lake	3 June 2017	3*	4	1.3
Mean					1.0
Amchitka	Infantry Road	9 June 2015	21	12	0.6
	Jones Lake-Engineer Road	9 June 2015	17	3	0.2
	Charlie to Baker Taxiway	9 June 2015	4	0	0.0
Mean					0.4
Attu	Old Loran/Old Runways	3 June 2015	4	0	0.0
	Massacre to Top Engineer	4 June 2015	7	0	0.0
	Navytown to Quonset Valley	4 June 2015	6	0	0.0
Mean					0.0

\*High winds did not allow completion or resurveys of three of six routes.

## Results

Rock Ptarmigan were heard or seen on all but one (2017 only) point-count routes on Adak and two of three on Amchitka but none was recorded on any of the three point-count routes on Attu (Table 2). However, one ptarmigan pair was seen and four males were heard prior to establishment of point-count routes but not near any of the point-count stops on Attu. Numbers of ptarmigan per stop recorded on point-count routes were highest (1.9, 1.4, 1.0; 2015–2017, respectively) on Adak, lower (0.4) on Amchitka, and non-existent (0.0) on Attu.

## Discussion

A literature review of surveys and reports of Rock Ptarmigan on Adak, Amchitka, and Attu Islands revealed that Rock Ptarmigan were mentioned but that few surveys occurred over time with the exception of Amchitka with less information for Attu and Adak (Appendix S1). Large populations were documented for Amchitka (White *et al.* 1977) and Attu (Braun *et al.* 2014) over short periods. Overall, the literature suggests populations of Rock Ptarmigan on the three islands were historically low, especially on Attu.

Our point counts indicate the Rock Ptarmigan population was highest on Adak (2015–2017), lower in 2015 on Amchitka, and very low on Attu in 2015. Our point-count survey data on Attu in 2015 con-

firmed the ongoing decline on this island reported by Braun *et al.* (2014) from an intensive survey area conducted in 2003–2009. No effort was made to quantify ptarmigan numbers on Attu at higher elevations but ptarmigan were common at lower elevations in 2003–2005 (Braun *et al.* 2014).

The areas that we surveyed on all three islands had similar relief (low marine and stream terraces), were highly disturbed in the mid 1940s and 1950s (Amchitka and Attu) to the late 1990s (Adak), but are now well vegetated with low to non-existent recent human occupation. The three islands have similar predator assemblages (no ground mammals except rats, but with eagles, falcons, gulls, jaegers, owls, and ravens), but we have no estimate of densities. We have no basis to expect that predators (Gilg *et al.* 2003; Therrien *et al.* 2014) affected ptarmigan numbers on the three islands in 2015. We also detected no evidence that male aggressive behaviour was a factor at the densities we observed (Mougeot *et al.* 2003). The possibility that herbivory (Sinclair *et al.* 1988) could affect populations of ptarmigan across islands at substantial distances from each other through plant compounds was considered but was deemed unlikely because of isolation, few deciduous shrubs, and distances involved.

We documented three different levels of abundance of Rock Ptarmigan on Adak (high), Amchitka

(lower), and Attu (very low) in 2015. The apparent, long-term decline on Attu since 2003 (Braun *et al.* 2014) appears to have stabilized from 2009 to 2014 (Braun *et al.* 2014). We agree with the hypothesis of Sandercock *et al.* (2005) that animal cycles in Arctic marine and terrestrial environments are most likely affected by latitudinal gradients in the north and altitudinal gradients elsewhere. The islands we studied are surrounded by the North Pacific Ocean and the Bering Sea and we worked at or below 300 m, thus the birds on these high latitude islands are mostly affected by the marine environment. We further agree that systematic surveys (Tesar *et al.* 2016) to detect trends in breeding populations (Nichols and Williams 2006) of different populations of Rock Ptarmigan are needed at least at 3–5 year intervals for both theoretical and practical reasons and should be able to detect population changes. It is possible that further translocations, similar to the one from Attu to Agattu in 2003–2005 (Kaler *et al.* 2010), may be considered to re-establish populations where they were extirpated by introduced Arctic Fox (*Vulpes lagopus*). Braun *et al.* (2014) documented the immediate recovery of ptarmigan after removal of Arctic Fox. But before such future translocations can be considered, a better survey protocol was needed to determine population status and trends of ptarmigan on these other islands. Knowing when ptarmigan populations may be ‘high’, especially if they cycle, also would be important so adequate numbers can be captured for immediate release on islands currently unoccupied by Rock Ptarmigan. This should reduce costs and improve chances for success of the transplants. Understanding fluctuations of Rock Ptarmigan populations, if they occur, is also important in the Arctic as the results from studies on islands may have relevance to ‘cycles’ and management of species of ptarmigan in mainland areas.

Point counts may be the most efficient and least expensive method to obtain standardized data (all birds seen and/or heard) for Rock Ptarmigan in areas with road or trail systems because large areas can be surveyed with few personnel. Early counts (May) should provide an opportunity to record more females than counts in early to mid June when females will be nesting. The three islands of Adak, Amchitka, and Attu each have different Rock Ptarmigan subspecies of conservation importance (Pruett *et al.* 2010) and their population dynamics deserve further study. There is a continuing need for population data to provide insight into whether cycles exist and their periodicity and synchronicity among islands.

### Author Contributions

Conceptualization: C.E.B., W.P.T., and S.M.E.; Funding Acquisition: S.M.E.; Investigation: C.E.B.,

W.P.T., S.M.E., and L.M.S.; Writing – Original Draft: C.E.B.; Writing – Review & Editing: C.E.B., W.P.T., S.M.E., and L.M.S.

### Acknowledgements

We thank the crew of U.S. Fish and Wildlife Service research ship *Tiglex* for safe transit along the North Pacific Ocean and the Bering Sea and especially the administrators of the Alaska Maritime National Wildlife Refuge for support of our efforts over a period of years. We acknowledge the help of the reviewers and the Editor for improvements in the manuscript. The findings and conclusions in this article are those of the author(s) and do not necessarily represent the views of the U.S. Fish and Wildlife Service.

### Literature Cited

- Amundsen, C.C., and E.E.C. Clebsch.** 1971. Dynamics of the terrestrial ecosystem vegetation of Amchitka Island, Alaska. *Bioscience* 21: 619–623. <https://doi.org/10.2307/1295734>
- AOU (American Ornithologists’ Union).** 1957. Checklist of North American birds. Fifth Edition. American Ornithologists’ Union, Washington, DC, USA.
- Armstrong, R.H.** 1971. Physical climatology of Amchitka Island, Alaska. *Bioscience* 21: 607–609. <https://doi.org/10.2307/1295730>
- Armstrong, R.H.** 1977. Weather and climate. Pages 53–58 in *The Environment of Amchitka Island*. Edited by M.L. Merritt and R.G. Fuller. U.S. Energy Research and Development Administration, TID-26712. U.S. Department of Energy, Washington, DC, USA.
- Bart, J., M. Fuller, P. Smith, and L. Dunn.** 2011. Use of large-scale, multi-species surveys to monitor Gyrfalcon and ptarmigan populations. Pages 263–271 in *Gyrfalcons and Ptarmigan in a Changing World*. Edited by R.T. Watson, T.J. Cade, M. Fuller, G. Hunt, and E. Potapov. The Peregrine Fund, Boise, Idaho, USA. Accessed 2 March 2018. <http://www.peregrinefund.org/subsites/conference-gyr/proceedings/>.
- Braun, C.E., W.P. Taylor, and S.M. Ebbert.** 2014. Changes in Evermann’s Rock Ptarmigan density on an eastern portion of Attu Island, Alaska, 2003–2009. *Northwestern Naturalist* 95: 28–34. <https://doi.org/10.1898/NWN13-10.1>
- Brodsky, L.M., and R.D. Montgomerie.** 1987. Asymmetrical contests in defence of Rock Ptarmigan territories. *Behavioral Ecology and Sociobiology* 21: 267–272. <https://doi.org/10.1007/bf00292508>
- Bystrak, D.** 1981. The North American breeding bird survey. *Studies in Avian Biology* 6: 34–41.
- Cattadori, I.M., D.T. Haydon, and P.J. Hudson.** 2005. Parasites and climate synchronize Red Grouse populations. *Nature* 433: 737–740. <https://doi.org/10.1038/nature03276>
- Coats, R.R.** 1956. Geology of northern Adak Island, Alaska. U.S. Department of the Interior, Geological Survey Bulletin 1028-C. U.S. Government Printing

- Office, Washington, DC, USA. Accessed 2 March 2018. <https://pubs.usgs.gov/bul/1028c/report.pdf>.
- Cotter, R.C.** 1991. Population attributes and reproductive biology of Rock Ptarmigan (*Lagopus mutus*) in the central Canadian Arctic. M.Sc. thesis, University of Alberta, Edmonton, Alberta, Canada.
- Cotter, R.C.** 1999. The reproductive ecology of Rock Ptarmigan (*Lagopus mutus*) in the central Canadian Arctic. *Arctic* 52: 23–32. <https://doi.org/10.14430/arctic.906>
- Dobson, A.P., and P.J. Hudson.** 1992. Regulation and stability of a free-living host–parasite system: *Trichostrongylus tenuis* in Red Grouse. II: Population models. *Journal of Animal Ecology* 61: 487–498. <https://doi.org/10.2307/5339>
- Droege, S.** 1990. Survey designs and statistical methods for the estimation of avian population trends. Pages 1–4 in *The North American Breeding Bird Survey*. Edited by J.R. Sauer and S. Droege. Biological Report 90 (1). U.S. Department of the Interior, Fish and Wildlife Service, Washington, DC, USA. Accessed 2 March 2018. <https://pubs.er.usgs.gov/publication/5230211>.
- Elton, C.S.** 1924. Periodic fluctuations in the numbers of animals: their causes and effects. *British Journal of Experimental Biology* 2: 119–163. Accessed 2 March 2018. <https://pdfs.semanticscholar.org/22fe/348adeffd92761fa5a5396a71c7820f5b8c7.pdf>.
- Everett, K.R.** 1971. The structure and origin of the organic soil cover of Amchitka Island, Alaska. *Bioscience* 21: 618. <https://doi.org/10.2307/1295733>
- Favaron, M., G.C. Scherini, D. Preatoni, G. Tosi, and L.A. Wauters.** 2006. Spacing behavior and habitat use of rock ptarmigan (*Lagopus mutus*) at low density in the Italian Alps. *Journal of Ornithology* 147: 618–628. <https://doi.org/10.1007/s10336-006-0087-z>
- Fraser, G.D., and G.L. Snyder.** 1959. Geology of southern Adak Island and Kagalaska Island, Alaska. U.S. Department of the Interior, Geological Survey Bulletin 1028-M. U.S. Government Printing Office, Washington, DC, USA. Accessed 2 March 2018. <https://pubs.usgs.gov/bul/1028m/report.pdf>.
- Gates, O., H.A. Powers, and R.E. Wilcox.** 1971. Geology of the Near Islands, Alaska. U.S. Department of the Interior, Geological Survey Bulletin 1028-U. U.S. Government Printing Office, Washington, DC, USA. Accessed 2 March 2018. <https://pubs.er.usgs.gov/publication/b1028U>.
- Gibson, D.D., and B. Kessel.** 1997. Inventory of the species and subspecies of Alaska birds. *Western Birds* 28: 45–95. Accessed 2 March 2018. [https://www.westernfieldornithologists.org/archive/V46/46\(2\)-p094-p185.pdf](https://www.westernfieldornithologists.org/archive/V46/46(2)-p094-p185.pdf).
- Gilg, O., I. Hanski, and B. Sittler.** 2003. Cyclic dynamics in a simple vertebrate predator-prey community. *Science* 302: 866–868. <https://doi.org/10.1126/science.1087509>
- Holder, K., and R. Montgomerie.** 1993. Rock Ptarmigan (*Lagopus mutus*). In *The Birds of North America*, No. 51. Edited by A. Poole and F. Gill. Academy of Natural Sciences, Philadelphia, Pennsylvania, USA.
- Hudson, P.J., D. Newborn, and A.P. Dobson.** 1992. Regulation and stability of a free-living host–parasite system: *Trichostrongylus tenuis* in Red Grouse. I: Monitoring and parasite reduction experiments. *Journal of Animal Ecology* 61: 477–486. <https://doi.org/10.2307/5339>
- Kaler, R.S.A., S.M. Ebbert, C.E. Braun, and B.K. Sandercock.** 2010. Demography of a reintroduced population of Evermann's Rock Ptarmigan in the Aleutian Islands. *Wilson Journal of Ornithology* 122: 1–14. <https://doi.org/10.1676/08-099.1>
- Magnússon, K., J. Brynjarsdóttir, and Ó.K. Nielson.** 2004. Population cycles in Rock Ptarmigan *Lagopus muta*: modelling and parameter estimation. RH-19-2004, Raunvísindastofnun Háskólans, Iceland.
- Montgomerie, R., and K. Holder.** 2008. Rock Ptarmigan (*Lagopus muta*). In *The Birds of North America*. Edited by A.F. Poole. Cornell Laboratory of Ornithology, Ithaca, New York, USA. <https://doi.org/10.2173/bna.51>
- Mougeot, F., S.M. Redpath, F. Leckie, and P.J. Hudson.** 2003. The effect of aggressiveness on the population dynamics of a territorial bird. *Nature* 421: 737–739. <https://doi.org/10.1038/nature01395>
- Nichols, J.D., and B.K. Williams.** 2006. Monitoring for conservation. *Trends in Ecology & Evolution* 21: 668–673. <https://doi.org/10.1016/j.tree.2006.08.007>
- Pedersen, A.O., B.-J. Bardsen, N.G. Yoccoz, N. Lecomte, and E. Fuglei.** 2012. Monitoring Svalbard Rock Ptarmigan: distance sampling and occupancy modeling. *Journal of Wildlife Management* 76: 308–316. <https://doi.org/10.1002/jwmg.276>
- Pedersen, A.O., O. Overrein, S. Unander, and E. Fuglei.** 2005. Svalbard Rock Ptarmigan (*Lagopus mutus hyperboreus*): a status report. Rapportserie No. 125, Norwegian Polar Institute, Tromsø, Norway. Accessed 2 March 2018. <http://hdl.handle.net/11250/173200>.
- Pelletier, L., and C.J. Krebs.** 1997. Line-transect sampling for estimating ptarmigan (*Lagopus* spp.) density. *Canadian Journal of Zoology* 75: 1185–1192. <https://doi.org/10.1139/z97-141>
- Powers, H.A., R.R. Coats, and W.H. Nelson.** 1960. Geology of and submarine physiology of Amchitka Island, Alaska. U.S. Department of Interior, Geological Survey Bulletin 1028-P. U.S. Government Printing Office, Washington, DC, USA. Accessed 2 March 2018. <https://pubs.usgs.gov/bul/1028p/report.pdf>.
- Pruett, C.L., T.N. Turner, C.M. Topp, S.V. Zagrebely, and K. Winker.** 2010. Divergence in an archipelago and its conservation consequences in Aleutian Island Rock Ptarmigan. *Conservation Genetics* 11: 241–248. <https://doi.org/10.1007/s10592-009-0026-7>
- Ralph, C.J., S. Droege, and J.R. Sauer.** 1995. Managing and monitoring birds using point counts: standards and applications. Pages 161–170 in *Monitoring Bird Populations by Point Counts*. Edited by C.J. Ralph, J.R. Sauer, and S. Droege. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station General Technical Report PSW-GTR-149. Albany, California, USA. Accessed 2 March 2018. [https://www.fs.fed.us/psw/publications/documents/gtr-149/pg161\\_168.pdf](https://www.fs.fed.us/psw/publications/documents/gtr-149/pg161_168.pdf).
- Robbins, C.S., D. Bystrak, and P.H. Geissler.** 1986. The Breeding Bird Survey: its first fifteen years, 1965–1979. U.S. Department of the Interior, Fish and Wildlife Service Resource Publication 157. Accessed 2 March 2018. <https://pubs.er.usgs.gov/publication/5230189>.



- Sandercock, B.K., K. Martin, and S.J. Hannon.** 2005. Life history strategies in extreme environments: comparative demography of Arctic and alpine ptarmigan. *Ecology* 86: 2176–2186. <https://doi.org/10.1890/04-0563>
- Sinclair, A.R.E., C.J. Krebs, J.N. Smith, and S. Boutin.** 1988. Population biology of snowshoe hares. III. Nutrition, plant secondary compounds and food limitation. *Journal of Animal Ecology* 57: 787–806. <https://doi.org/10.2307/5093>
- Taylor, W.P.** 2013. Status of upland game within Alaska's highway system: a comprehensive report focusing on 2007–2011. Wildlife Management Report, ADF&G/DWC/WMR-2013-1. Alaska Department of Fish and Game, Division of Wildlife Conservation, Palmer, Alaska, USA. Accessed 12 August 2015. [https://www.adfg.alaska.gov/static/research/programs/smallgame/pdfs/upland\\_game\\_status\\_2007\\_2011.pdf](https://www.adfg.alaska.gov/static/research/programs/smallgame/pdfs/upland_game_status_2007_2011.pdf)
- Tesar, C., M.-A. Dubois, and A. Shestakov.** 2016. Toward strategic, coherent, policy-relevant Arctic science. *Science* 353 (6306): 1368–1370. <https://doi.org/10.1126/science.aai8198>
- Therrien, J.F., G. Gauthier, E. Korpimäki, and J. Bêty.** 2014. Predation pressure by avian predators suggests summer limitation of small-mammal populations in the Canadian Arctic. *Ecology* 95: 56–67. <https://doi.org/10.1890/13-0458.1>
- Watson, A.** 1965. A population study of ptarmigan (*Lagopus mutus*) in Scotland. *Journal of Animal Ecology* 34: 135–172. <https://doi.org/10.2307/2373>
- Watson A., R. Moss, and P. Rothery.** 2000. Weather and synchrony in 10-year population cycles of Rock Ptarmigan and Red Grouse in Scotland. *Ecology* 81: 2126–2136. [https://doi.org/10.1890/0012-9658\(2000\)081\[2126:wasiyp\]2.0.co;2](https://doi.org/10.1890/0012-9658(2000)081[2126:wasiyp]2.0.co;2)
- Weatherbase.** 2015. Travel weather averages for Adak, Alaska, USA. Accessed 12 August 2015. <http://www.weatherbase.com/adak>
- Weeden, R.B.** 1965. Breeding density, reproductive success and mortality of Rock Ptarmigan at Eagle Creek, central Alaska from 1960–1964. *Transactions of the North American Wildlife Conference* 30: 336–348.
- White, C.M., F.S.L. Williamson, and W.B. Emison.** 1977. Avifaunal investigations. Pages 227–260 in *The Environment of Amchitka Island*. Edited by M.L. Merritt and R.G. Fuller. U.S. Energy Research and Development Administration, TID-26712. U.S. Department of Energy, Washington, DC, USA.

Received 3 May 2017

Accepted 6 February 2018

### Supplementary Material:

**Appendix S1.** Historical review of information and previous studies and surveys for Rock Ptarmigan (*Lagopus muta*) on three islands (Adak, Amchitka, and Attu) in the Western Aleutian Islands in Alaska.

# The Canadian Field-Naturalist

## SUPPLEMENTARY MATERIAL:

### Monitoring Rock Ptarmigan (*Lagopus muta*) populations in the Western Aleutian Islands, Alaska Clait E. Braun, William P. Taylor, Steven M. Ebbert, and Lisa M. Spittler

**Appendix S1.** Historical review of information and previous studies and surveys for Rock Ptarmigan (*Lagopus muta*) on three islands (Adak, Amchitka, and Attu) in the Western Aleutian Islands in Alaska.

All islands studied had been disturbed by historical military activities. The military presence on Amchitka ended in 1951 (Kenyon 1961), the Naval Air Facility closed on Adak in 1997, and in 2010 the small Coast Guard station on Attu was closed. Hunting of Rock Ptarmigan on Adak continues but possibly at lower levels than when the military began downsizing in 1994. Despite the extensive military presence, the published literature on numbers of Rock Ptarmigan in the Aleutian Islands is limited and highly fragmented over time. Sutton, Taber, and Wilson were military officers stationed in the Aleutian Islands during World War II but were also trained biologists / ornithologists and published in ornithological journals. Standardized surveys conducted at intervals are limited for all three islands. The few references, outside of that on Amchitka, suggest that ptarmigan numbers were low with few exceptions.

#### Adak Island

Taber (1946) was stationed on Adak and reported on his bird observations during 18 November 1945 to 20 January 1946. He does not report any ptarmigan during this period even though his observation locales on the northern tip of the island included Andrew Lake (“Lagoon”) and Clam Lagoon. It is possible that activities of armed forces personnel could have influenced the abundance of ptarmigan in this area although Taber (1946: 273) speculated “the long-range effects [of military activities] upon the bird populations to have been small”. He also compared (pp. 276–277) the winter avifaunas of Adak and Attu based on Sutton and Wilson (1946) without mention of Rock Ptarmigan. Despite the military activity on Adak during the 1940s through the 1990s, little information is available on the presence of Rock Ptarmigan on this island other than the early (1907) description of the subspecies (*atkhensis*) by Clark (AOU 1957: 134). The only data found on population numbers (based on intensive field surveys by four observers) on Adak were those of Masteller (1981) who reported 3.0 males/km<sup>2</sup> on a 11.9 km<sup>2</sup> area during 13–15 May 1981 in the Mt. Adagdak re-

gion. Audubon Christmas Bird Count data (National Audubon Society 2015; <http://netapp.audubon.org/cbcobservation/>) reported for Adak for 1973 through 1994 revealed that numbers of observers, ptarmigan observed, and time afield fluctuated without apparent patterns.

#### Amchitka Island

Substantial information is available on Rock Ptarmigan on Amchitka, especially food habits and body mass (Elison 1980; Emison and White 1988), and plumage moult (Jacobsen *et al.* 1983). These studies were part of a wide-ranging environmental data collection program funded by the U.S. Atomic Energy Commission (AEC) and summarized by Amundsen and Clebsch (1971) and Armstrong (1971). The first listing of birds on Amchitka was by Kenyon (1961) from data collected in 1955–1959 following the departure of the military in 1951. His field investigations were after the initial intensive removal program for Arctic Fox (*Vulpes lagopus*) which began in 1950 (Jones 1960; Kenyon 1961; Emison *et al.* 1985). It appears most of the foxes had been removed from the eastern portion of Amchitka by about 1954, but some were still common on the western (less frequently visited) portion until at least 1955. Large drops of pesticide-treated baits in 1956 and 1957 probably reduced the fox population to only a few animals (and no active dens), which eventually died out sometime between 1962 and 1965 (Thomson and Byrd 2010). Thus, Kenyon’s field work overlapped several years of intensive predator removal. The only land mammal present after the eradication program were rats (*Rattus* spp.). Kenyon (1961: 318) indicates Rock Ptarmigan were “quite scarce” during 1955–1957 and “much more abundant in 1959, perhaps because of the reduction of foxes”. Only two to four ptarmigan were seen in a 22-km drive prior to 1959 whereas 15–20 birds were observed on this same route in 1959. White *et al.* (1977) initiated field work on Amchitka as part of the environmental data collection program funded through the AEC. Their field work continued from August 1967 through September 1973 and included

all months of the year. They surveyed a 1012-ha plot for the presence of Rock Ptarmigan and recorded 32 pairs in 1969 and 34 pairs in 1970. No surveys were conducted in 1971, but their observations indicated the size of the breeding population was similar to those recorded in 1969–1970. White *et al.* (1977: 242) estimated the spring and summer population size, based on extrapolation from their intensive study area, was 1400 Rock Ptarmigan whereas the autumn population was 2800, and the winter population was 2500 birds for the entire island. We do not believe the birds were moving on and off the island seasonally and the White *et al.* (1977) estimates from their intensive surveys on their study block were used by them to illustrate annual fluctuations and to estimate population sizes.

Zeillemaker and Schulmeister (1984: 5) made a roadside survey along Infantry Road between “mile post 1 and mile post 22” on Amchitka Island on 1 May 1984 and reported 34 Rock Ptarmigan. More recently, Scharf (1994) conducted surveys on Amchitka and indicated Rock Ptarmigan had increased over the levels reported in the late 1960s. This finding was based on surveys during 7 and 10 June 1994 using six observers “walking abreast” through four subplots (each 50 ha with nine pairs found in 1994 in 200 ha) within the 1012-ha plot described by White *et al.* (1977). Driving surveys were also established in 1994 along 8.3 km of Infantry Road and a route of similar length on Engineer Road starting at the east end of Baker Runway. A point-count survey with 12 survey points spaced at 400 m intervals was also established in 1979. The vehicle surveys recorded eight Rock Ptarmigan in 1994, 13 in 1988, six in 1980, and six in 1979 (Scharf 1994: Table 6). All of these surveys were conducted between 1 and 22 June.

#### Attu Island

The early reports (1910, 1925 cited in Bent 1932: 231) of Rock Ptarmigan on Attu suggested this species occurred at low densities in the higher mountains. Only three birds were reported in one day afield in 1910 and three in three days in 1925. Sutton and Wilson (1946) and Wilson (1948) were stationed on Attu for short periods (Sutton from 20 February to 18 March 1945, Wilson from September 1944 to October 1945). Sutton and Wilson (1946: 83) indicated they did not personally see ptarmigan despite making “a point of looking closely for them”. They report second-hand accounts of two birds in full winter plumage 3.2 km west of the head of Massacre Bay on 7 March. Both authors were on Attu when it was an active military base and spent most of their time afield from Casco Cove and Murder Point to Massacre Bay with short trips to Alexi Point, Temnac Bay, and

Chirikof Point. Wilson (1948: 127) suggested Rock Ptarmigan were restricted to the “rugged mountain slopes of the interior in summer” on Attu and that he never saw “it or its tracks anywhere along the coast, even in the dead of winter”. He reports a male on 10 June 1945 on Fishhook Ridge, a pair on 12 July on the west shoulder of Terrible Mountain, and a male and two females on 28 July high on the steep mountain southwest of Casco Cove. All of the observations in the mid-1940s were during the active military presence and prior to when Arctic Fox had been removed (1999–2000) from the island.

More recently, Haflinger and Tobish (1977) conducted a population survey for Rock Ptarmigan along walking transects on Attu in the southeast portion of the island from Casco Cove to Massacre Bay and inland to the base of the mountains. They reported densities of 2.4 pairs/km<sup>2</sup> for breeding ptarmigan on Attu in 1977. Holder (1994) estimated there were 0.9 territories/km<sup>2</sup> for Rock Ptarmigan on her 10 km<sup>2</sup> study area on Attu in 1994. Her data indicated that territories were primarily on the higher slopes of the island. These densities are lower than those reported for 2003 (4 pairs/km<sup>2</sup>) by Braun *et al.* (2014) prior to a transplant of ptarmigan from Attu to Agattu (Kaler *et al.* 2010).

Introduced Arctic Fox were removed from Amchitka between 1950 and 1965 (Thomson and Byrd 2010), Attu between 1999 and 2000 (Ebbert and Byrd 2002), and from Adak between 2002 (Scharf *et al.* 2002) and 2004 (Alaska Maritime National Wildlife Refuge, Homer, Alaska, USA unpubl. data).

#### Literature Cited

- Bent, A.C.** 1932. Life histories of North American gallinaceous birds. U.S. National Museum Bulletin 162. <https://doi.org/10.5479/si.03629236.162.i>
- Ebbert, S.M., and G.V. Byrd.** 2002. Eradications of invasive species to restore natural biological diversity on Alaska Maritime National Wildlife Refuge. Pages 102–109 in *Turning the Tide: the Eradication of Invasive Species*. Edited by C.R. Veitch and M.N. Clout. Invasive Specialist Group, Gland, Switzerland and Cambridge, United Kingdom. Accessed 2 March 2018. [http://www.issg.org/pdf/publications/turning\\_the\\_tide.pdf](http://www.issg.org/pdf/publications/turning_the_tide.pdf).
- Elison, G.W.** 1980. Food habits study of Rock Ptarmigan (*Lagopus mutus*) on Amchitka Island, Alaska. M.Sc. thesis, Washington State University, Pullman, Washington, USA.
- Emison, W.B., and C.M. White.** 1988. Foods and weights of the Rock Ptarmigan on Amchitka, Aleutian Islands, Alaska. *Great Basin Naturalist* 48: 533–540.
- Emison, W.B., C.M. White, and F.S.L. Williamson.** 1985. Life history and foods of the Rock Ptarmigan on Amchitka Island, Alaska with comments on factors influencing population fluctuations. Unpublished Report 85/30. Aleutian Islands Unit, Alaska Maritime National

- Wildlife Refuge, Homer, Alaska, USA.
- Haflinger, K., and T. Tobish.** 1977. Results of a population survey of the Rock Ptarmigan *Lagopus mutus evermanni* on Attu Island, Alaska, May 12–August 5 1977. Unpublished Report. U.S. Department of the Interior, Fish and Wildlife Service, Alaska Maritime National Wildlife Refuge, Homer, Alaska, USA.
- Holder, K.** 1994. Evolutionary divergence of Rock Ptarmigan: an Aleutian solution. *Information North*. Arctic Institute of North America 20(4): 1–4.
- Jacobsen, E.E., C.M. White, and W.B. Emison.** 1983. Molting adaptations of Rock Ptarmigan on Amchitka Island, Alaska. *Condor* 85: 420–426. <https://doi.org/10.2307/1367980>
- Jones, R.D.** 1960. Refuge Narrative Report April thru August, 1960. Aleutian Islands National Wildlife Refuge, Bureau of Sport Fisheries and Wildlife, Cold Bay, Alaska, USA.
- Kenyon, K.W.** 1961. Birds of Amchitka Island, Alaska. *Auk* 78: 305–326. <https://doi.org/10.2307/4082270>
- Masteller, M.** 1981. Ptarmigan census results, Adak Island, May 1981. Unpublished Report. Alaska Maritime National Wildlife Refuge, Aleutians Islands Unit, Homer, Alaska, USA.
- National Audubon Society.** 2015. Christmas Bird Counts. National Audubon Society, New York. Accessed 12 August 2015. <http://netapp.audubon.org/cbcobservation/>
- Scharf, L.** 1994. Aleutian Canada Goose surveys and other biological surveys on Amchitka Island, Alaska, 31 May–18 June 1994. U.S. Fish and Wildlife Service, Alaska Maritime National Wildlife Refuge, Aleutians Islands Unit, Adak, Alaska, USA.
- Scharf, L., G. Thomson, and S. Ebbert.** 2002. Eradication of Arctic foxes at Adak Island, Alaska in 2002. U.S. Department of the Interior, Fish and Wildlife Service Report. Alaska Marine National Wildlife Refuge 02/02. Homer, Alaska, USA.
- Sutton, G.M., and R.S. Wilson.** 1946. Notes on the winter birds of Attu. *Condor* 48: 83–91. <https://doi.org/10.2307/1363994>
- Taber, R.D.** 1946. Winter birds of Adak, Alaska. *Condor* 48: 272–277. <https://doi.org/10.2307/1364080>
- Thomson, G., and V. Byrd.** 2010. Summary of eradication efforts of Arctic foxes at Amchitka Island, Alaska from 1949 to 1965. U.S. Department of the Interior, Fish and Wildlife Service Report. Alaska Marine National Wildlife Refuge 2010/09. Homer, Alaska, USA.
- White, C.M., F.S.L. Williamson, and W.B. Emison.** 1977. Avifaunal investigations. Pages 227–260 in *The Environment of Amchitka Island*. Edited by M.L. Merritt, and R.G. Fuller. U.S. Energy Research and Development Administration, TID-26712. U.S. Department of Energy, Washington, DC, USA.
- Wilson, R.S.** 1948. The summer bird life of Attu. *Condor* 50: 124–129. <https://doi.org/10.2307/1364686>
- Zeilemaker, C.F., and R. Schulmeister.** 1984. Bird and mammal observations at Amchitka Island, Aleutian Islands, Alaska 30 April–10 May 1984. BIR-MAM4 (Bird-Mammal Disk). U.S. Department of the Interior, Fish and Wildlife Service, Aleutian Islands Unit, Alaska Maritime National Wildlife Refuge, Seattle, Washington, USA.